

Updated checklist of *Poa* in the Iberian Peninsula and Balearic Islands

Ana Ortega-Olivencia¹, Juan A. Devesa²

1 Área de Botánica, Facultad de Ciencias, Universidad de Extremadura, Avenida de Elvas, s.n., 06006 Badajoz, Spain **2** Departamento de Botánica, Ecología y Fisiología Vegetal, Facultad de Ciencias, Universidad de Córdoba, Campus de Rabanales, Edificio José Celestino Mutis, Ctra. de Madrid km. 396 A, 14014 Córdoba, Spain

Corresponding author: Ana Ortega-Olivencia (aortega@unex.es)

Academic editor: M. Nobis | Received 20 April 2018 | Accepted 27 June 2018 | Published 10 July 2018

Citation: Ortega-Olivencia A, Devesa JA (2018) Updated checklist of *Poa* in the Iberian Peninsula and Balearic Islands. *PhytoKeys* 103: 27–60. <https://doi.org/10.3897/phytokeys.103.26029>

Abstract

Based on our study of 4,845 herbarium sheets of the genus *Poa* from the area covered by *Flora iberica*, namely, the Iberian Peninsula and the Balearic Islands, we recognise 24 taxa (17 species, 1 subspecies and 8 varieties), mostly perennials. Most of these taxa have wide global and/or European distributions, while two (*P. legionensis* and *P. minor* subsp. *nevadensis*) are Spanish endemics and two have restricted distributions (*P. ligulata*, Iberia–North Africa; *P. flaccidula*, Iberia–North Africa and the Balearic Islands, extending to Provence, France). We have studied the original publications of more than 225 names considered as synonyms, with those more historically cited in *Flora iberica* taken into account in this paper; a total of 26 are new synonyms. The following names are typified: *P. alpina* var. *involutrata* Lange, *P. annua* var. *lanuginosa* Sennen, *P. minor* subsp. *nevadensis* Nannf., *P. paui* Font Quer, *P. sulcata* Lag. and *P. trivialis* var. *flaccida* Willk. ex J.J. Rodr. We include *P. compressa* L. in the flora of Portugal for the first time and present detailed illustrations of three very interesting taxa (*P. legionensis*, *P. minor* subsp. *nevadensis* and *P. ligulata*). In addition to a general species key, we provide the following information for each taxon: synonyms, types, typification, the most relevant iconography, regional flowering time, regional and general distribution and, as supplementary material, the number of sheets examined and a list of selected materials.

Keywords

Checklist, *Flora iberica*, Gramineae, Poaceae, *Poa*, Portugal, Spain, taxonomy, typification

Introduction

The genus *Poa* L., included within subfamily Pooideae, supertribe Poodae and subtribe Poinae (Soreng et al. 2015), is considered to be monophyletic. This monophyly is supported by analyses of plastid and nuclear DNA markers (Gillespie et al. 2008). In addition, evidence for reticulation between this genus and other genera in Poinae has been uncovered; the same is true within the genus itself (e.g. *P. annua*) (Soreng et al. 2010). The genus comprises approximately 550 annual and perennial species (Soreng et al. 2017) of cosmopolitan distribution, primarily in cold and temperate regions. Most species are polyploids, but 9% are diploids, with an additional 4%–6%, mostly in Europe and rarely in Asia, having both diploid and polyploid populations (Soreng et al. 2010, Giussani et al. 2016). Many species are important weeds (e.g. *P. annua*), while others are cultivated for forage (e.g. *P. pratensis*) or used in pastures (e.g. *P. trivialis*, *P. alpina* and *P. bulbosa*) or lawns and golf courses (e.g. *P. nemoralis* and *P. pratensis*) (Watson and Dallwitz 1992).

The genus is characterised by a great diversity of sexual systems and its species can be strictly hermaphroditic, the most common reproductive system, or diclinous (Giussani et al. 2016). Apomictic reproduction by seeds, either facultative or obligate, is common in some species (reviewed in Soreng and Peterson 2012). The production of pseudoviviparous/bulbiferous spikelets occurs in some species, such as *P. bulbosa* and, to a lesser extent, *P. alpina*.

The first taxonomic treatment of *Poa* on the Iberian Peninsula, by Willkomm (1870), was remarkable: he recognised 15 species and numerous varieties in the territory of Spain and later increased this total by two (Willkomm 1893). In the 20th century, 19 of the 53 species and subspecies recognised by Edmonson (1980) in Europe were included in *Flora iberica*, a publication also encompassing the Balearic Islands. Of these, slightly less than half were listed for Portugal (see also Franco and Rocha 1998). The latest revision to *Poa* on the peninsula was carried out by Hernández Cardona (1978), who basically followed Edmonson. In this revision, Hernández Cardona also defined a new section (*Flaccidula* Á.M. Hern.) to accommodate *P. flaccidula* and recognised 19 species and subspecies plus two varieties.

In a recently completed revision of the genus *Poa* for a future volume (XIX) of *Flora iberica*, we recognised 18 species and subspecies and 8 varieties. The main aim of the present paper was to present an updated checklist of the genus. The information provided includes a general key to accepted taxa as well as a list of their most important synonyms, many of which are unknown outside of the Iberian Peninsula because they are found only on herbarium sheets or published in works of limited distribution. Some of the synonyms and an accepted name are typified and updated information on the ecology and flowering characteristics of each taxon in the covered territory is given along with its regional and worldwide distribution.

Methods

The taxonomic classification scheme followed in this paper, which begins with the type species *P. pratensis*, reflects currently understood relationships amongst recognised sections in the genus. An infrageneric classification of accepted species of *Poa* in the Iberian flora is also presented in the Results section.

We reviewed 4,845 sheets housed in the following herbaria: BC, BC-Sennen, C-Lange, COI, COI-Willk., G-Boiss., GDA-GDAC, HGM, HSS, JACA, MA, MAF, MGC, SALA-SALAF, SEV, UPP-Nannf. and UNEX (acronyms according to Thiers, continuously updated). We studied the most important synonyms of each accepted name and consulted the original publications, with a special focus on names directly related to the territory covered by *Flora iberica*. For each accepted taxon, we recorded synonyms, types (type protologue) and, in some cases, the typification. After studying the herbarium sheets, we obtained updated information on flowering phenology and the ecology of the area. We also researched the worldwide distribution of each taxon and its presence or absence in each province covered by *Flora iberica*, including the territories of Andorra (And.), Portugal (Port.) and continental Spain (Spa.) plus the Balearic Islands. In the taxonomic treatment that follows, those provinces are ordered alphabetically using the same abbreviations given in *Flora iberica* (<http://www.floraiberica.es/>; see Fig. 1). If the name of a province appears in parentheses, a bibliographic citation exists but no herbarium sheet was studied to confirm it, while a question mark indicates that



Figure 1. Map of distribution of the provinces covered by *Flora iberica* (<http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=476&Pagina=27>).

the bibliographic citation is not entirely reliable. A selected list of herbarium sheets studied from each province is provided in the Suppl. material 1. Finally, some observations are included as explanatory notes for most species and subspecies.

Results and discussion

Currently, a total of 24 taxa are recognised: 17 species, 1 subspecies and 8 varieties. All are perennials except for *P. annua* and *P. infirma*, which are annuals.

Key to the species of *Poa* in Flora iberica

- 1 Plant annual, sometimes multicaulous **2**
- Plant perennial, cespitose, rhizomatous and/or stoloniferous..... **3**
- 2 Spikelets (1.6–)4–7.7(–9) mm; anthers (0.6–)0.7–1.3 mm, much longer than wide; caryopsis 1.3–2.1 mm **12. *P. annua***
- Spikelets 2.8–4.8 mm; anthers 0.2–0.4(–0.6) mm, the same as or slightly longer than the width; caryopsis 1–1.4 mm **13. *P. infirma***
- 3 Plant cespitose, without rhizomes and almost always without stolons, usually with basal bulbils and/or stems with thickened bases covered by old sheaths, these often split in fibres **4**
- Plant cespitose, with rhizomes and/or stolons, lacking bulbils or strongly thickened stem bases surrounded by old sheaths or, if present, then with the ligule of basal leaves 0.15–0.4(–0.8) mm, truncate..... **9**
- 4 Lemma with 5 prominent veins and base very hairy, with hairs longer than the width of the lemma..... **1. *P. pratensis***
- Lemma with 5 inconspicuous veins and base glabrous or with hairs much shorter than the width of the lemma..... **5**
- 5 Plant without stolons, bulbils or stems thickened at the base, glaucous; glumes lanceolate..... **5. *P. glauca***
- Plant with short stolons rooting at the nodes or with bulbils or thickened stem bases covered by old sheaths, green or glaucous; glumes lanceolate or the lowermost one narrowly subulate..... **6**
- 6 Plant cespitose, rarely with short stolons rooting at the nodes, without basal bulbils or thickened stem bases; lower glume narrowly subulate..... **6. *P. nemoralis***
- Plant cespitose, with basal bulbils and/or strongly thickened stem bases covered by old fibrous sheaths; lower glume lanceolate or ovate-lanceolate..... **7**
- 7 Basal and shoot leaves with ligule 0.3–2(–3.8) mm, the oldest 0.3–0.7 mm, ± truncate, blade (0.7–)1.5–4.5(–7.5) mm wide, flat **17. *P. alpina***
- Basal and shoot leaves with ligule 2–10.5 mm, ± oblong or triangular, often lacinate in acute triangular segments, blade 0.4–3.5(–4) mm wide, flat, conduplicate or convolute **8**

- 8 Plant with basal bulbils, frequently also present in the inflorescence; basal and shoot leaves filiform or linear, less frequently lanceolate; ligule membranous-hyaline; tuft concoloured, green or brown..... **15. *P. bulbosa***
- Plant without basal bulbils or bulbils in the inflorescence; basal and shoot leaves linear or lanceolate; ligule pearly white; tuft bicoloured, green and white because of the brightness of the ligules **16. *P. ligulata***
- 9 Branches of the inflorescence and/or spikelet peduncles smooth or nearly so, glabrous **10**
- Branches of the inflorescence and/or spikelet peduncles antrorse-scabrid... **14**
- 10 Ligule of the basal leaves 0.15–0.7(–0.8) mm, truncate **11**
- Ligule of the basal leaves 0.6–3.4(–4.3) mm, more or less ovate or triangular-ovate, not truncate **12**
- 11 Upper leaves with ligule 0.3–0.6(–0.8) mm, truncate-dentate; lemma base with hairs longer than the width of the lemma; plant cespitose-rhizomatous. **2. *P. legionensis***
- Upper leaves with ligule 0.7–2.5(–4.7) mm, more or less truncate or ovate, sometimes split into 2 or several parts; lemma base glabrous; plant densely cespitose **17. *P. alpina***
- 12 Palea with keels appressed-hairy or ciliate, rarely smooth and glabrous; anthers (1.2–)1.6–2.1 mm; branches of the inflorescence patent or reflexed after anthesis; upper flower of the spikelet female..... **14. *P. supina***
- Palea with keels antrorse-scabrid, never appressed-hairy or ciliate; anthers 0.7–1.7 mm; branches of the inflorescence erect or erect-patent; upper flower of the spikelet hermaphroditic **13**
- 13 Anthers 0.7–1.2 mm; branches of the inflorescence 0.15–0.2 mm in diameter, sulcate, \pm rigid, glabrous and smooth; spikelets with peduncle glabrous, smooth, having 2–4 flowers **8. *P. laxa***
- Anthers 0.8–1.7 mm; branches of the inflorescence c. 0.05–0.15 mm in diameter, not sulcate, \pm flexuous, glabrous and smooth, sometimes very loosely antrorse-scabrid; spikelets with peduncle glabrous or sometimes laxly antrorse-scabrid, having 4–7 flowers..... **9. *P. minor***
- 14 Stems compressed **15**
- Stems not or only slightly compressed..... **21**
- 15 Lemma and palea with hairy surfaces between veins **11. *P. flaccidula***
- Lemma and palea with glabrous surfaces between veins **16**
- 16 Ligule of the upper leaf lanceolate, generally acute, longer than the width of the blade **10. *P. trivialis***
- Ligule of the upper leaf truncate, obtuse, shorter or subequal to the width of the blade **17**
- 17 Lemma with conspicuous veins..... **18**
- Lemma with inconspicuous veins..... **19**
- 18 Stems with base 2.5–9 mm wide, usually very compressed; lemma with 5 glabrous veins; blade of the basal leaves (3.5–)5.5–11.2 mm wide, lanceolate

- or oblong-lanceolate **4. *P. chaixii***
- Stems with base 1.5–2.5(–3.5) mm wide, slightly compressed; lemma with 5 veins, the central and marginal ones appressed-hairy; blade of the basal leaves 0.5–3.8 mm wide, lanceolate, linear or setaceous **1. *P. pratensis***
- 19 Usually with two keels in the stems; glumes lanceolate or ovate-lanceolate, ± convergent; spikelet with rachilla usually glabrous; inflorescence with 1–2(–5) branches in the basal node, short and appressed **7. *P. compressa***
- Without keels in the stems; glumes subulate or lanceolate-subulate, straight and ± divergent; spikelet with rachilla glabrous or pubescent; inflorescence with 2–5 branches in the basal node, short or long, erect, rarely erect-patent or non-appressed on the axis **20**
- 20 Plant rhizomatous; leaves of the shoots usually distichous; spikelets 4.5–7(–7.6) mm; lower glume (2.8–)3.2–4.2 mm, lanceolate **3. *P. cenisia***
- Plant without rhizomes, sometimes with short stolons; leaves of the shoots non-distichous, sparse; spikelets 3–4.8(–5.5) mm; lower glume 2–3.5 mm, subulate **6. *P. nemoralis***
- 21 Lemma and palea with surfaces between veins hairy **11. *P. flaccidula***
- Lemma and palea with surfaces between veins glabrous **22**
- 22 Ligule of the upper leaf usually lanceolate, acute or acuminate, longer than the width of the blade; stems usually retrorse-scabrid around the nodes; plant stoloniferous; callus of lemma with hairs longer than the width of the lemma, very rarely glabrous **10. *P. trivialis***
- Ligule of the upper leaf ovate or oblong, obtuse, truncate or dentate, much shorter or subequal to the width of the blade; stems smooth around the nodes; plant rhizomatous, rarely with stolons; callus of lemma without hairs or with hairs larger or smaller than the width of the lemma **23**
- 23 Plant caespitose, sometimes with short stolons; glumes subulate or lanceolate-subulate, glossy; lemma with base glabrous or with hairs of much shorter length than the width of the lemma; ligule of the upper leaves 0.2–0.6(–0.8) mm **6. *P. nemoralis***
- Plant rhizomatous; glumes lanceolate, dull; lemma with base hairy, with hairs shorter or longer than the width of the lemma; ligule of the upper leaves usually 1–3.1 mm **24**
- 24 Plant ± glaucous; shoots usually with leaves distichous; culms usually scabrous below nodes; glumes subequal, ± straight or slightly converging with each other, 3-veined; lemma with 5 inconspicuous veins, with hairs at the base shorter or longer than the width of the lemma; palea with keels antrorsely-scaberulous or ciliolate, often something curly in the basal area **3. *P. cenisia***
- Plant usually green; shoots usually with non-distichous leaves; culms smooth below nodes; glumes unequal, curved or converging with each other, the lower with 1 or 3 veins; lemma with 5 conspicuous veins, with hairs at the base longer than the width of the lemma; palea with keels antrorsely-scaberulous... **1. *P. pratensis***

Checklist of *Poa* in the Iberian Flora

The species of *Poa* present in *Flora iberica* are classified into three subgenera and nine sections as indicated below. These taxonomic placements are provisional because some taxa (indicated by *) have not yet been subjected to DNA sequencing:

Poa subgen. *Poa* supersect. *Poa* sect. *Poa*: *P. pratensis*, **P. legionensis*, *P. cenisia*; supersect. *Homalopoa* (Dumort.) Soreng & L.J. Gillespie sect. *Homalopoa*: *P. chaixii*.

Poa subgen. *Stenopoa* (Dumort.) Soreng & L.J. Gillespie sect. *Stenopoa*: *P. glauca*, *P. nemoralis*; sect. *Tichopoa* Asch. & Graebn.: *P. compressa*; sect. *Oreinos* Asch. & Graebn.: *P. laxa*, *P. minor*; sect. *Pandemos* Asch. & Graebn.: *P. trivialis*; *sect. *Flaccidula* Á.M. Hern.: **P. flaccidula*.

Poa subgen. *Ochlopoa* (Asch. & Graebn.) Hyl. sect. *Micrantherae* Stapf: *P. annua*, *P. infirma*, *P. supina*; sect. *Arenariae* Stapf: *P. bulbosa*; sect. *Alpinae* (Nyman) Stapf: *P. ligulata*, *P. alpina*.

The sequence of species in this checklist is not alphabetical, but instead starts with sect. *Poa* because that section includes the type species; species are then ordered according to phylogenetic relationships, an arrangement more or less the inverse of that adopted by other authors, i.e. from more derived clades to those in a more basal position (see Gillespie et al. 2008; Soreng et al. 2010, 2017).

1. *Poa pratensis* L., Sp. Pl. 67. 1753 subsp. *pratensis*

Poa angustifolia var. *pratensis* (L.) Simonkai, Enum. Fl. Transsilv. 580. 1886.

Paneion pratense (L.) Lunell, Amer. Midl. Nat. 4: 222. 1915.

III. var. *pratensis* [Soreng and Peterson (2012: 67, fig. 18C–J, sub *P. pratensis* subsp. *pratensis*); Devesa (1987: 261, sub *P. pratensis*); var. *minor* [Soreng and Peterson (2012: 67, fig. 18 A–B, sub *P. pratensis* subsp. *irrigata*); var. *angustifolia* [Ruiz (1991: 29, lam. II, sub *P. angustifolia*); Soreng and Peterson (2012: 66, fig. 17H–I, sub *P. pratensis* subsp. *angustifolia*)].

Type. “Habitat in Europae pratis fertilissimis”. Typus: Russia, Prov. Sanct-Petersburg, 5 km australi-occidentum, versus a st. viae ferr. Mga. pratulum ad ripam dextram fl. Mga, 26 Jun 1997, N. N. Tzvelev N-257 (type conserved, designated by Soreng and Barrie 1999, pg. 157: BM-000576302; isoelectotypes: B, C, CAN, CONC, H, K, KW, L, LE, LIV, MA, MO, MW, NSW, P, PE, PR, S, SI, TNS, US, W).

Flowering. April–August (September).

Ecology. Grasslands at edges of watercourses, ravines, ponds and alpine wetlands (“borreguiles”), walls, wet soils on slopes, ditches, cultivated fields, clearings surrounded by pines, holm oaks, Portuguese oaks and other oaks; edaphically indifferent; 0–2400 m a.s.l.

Distribution. Eurasia, N Africa and Macaronesia (Azores, Madeira and Canary Islands); introduced in N, C and S America and Australia. Scattered throughout much of the Iberian Peninsula and Balearic Islands. **And. Spa.:** A Ab Al Av B Bu C Cc Co CR Cs Cu Ge Gr Gu Hu J L Le Lo Lu M Ma Mu Na O Or P PM[MI] (Po) S Sa Sg So SS (T) Te To V Va Vi Z Za. **Port:** AAl (BA) (BB) (BL) DL (E) Mi TM.

Notes. *Poa pratensis* is one of the most polymorphic taxa in the genus for a variety of reasons: its great morphological and cytological variation, the predominance of agamospermy, its vegetative propagation and wide distribution, the latter due in part to its introduction into many parts of the world for use on lawns, as fodder or for soil stabilisation (Soreng and Barrie 1999). At least 220 crop varieties are recognised (Stoneberg Holt et al. 2004).

In the territory covered by *Flora iberica*, three patterns of variation are recognised. Plants with scarcely any extravaginal shoots and possessing basal-leaf ligules with scattered or sometimes entangled apical hairs up to 0.4 mm and 0.2–0.5 mm on the back correspond to *Poa pratensis* var. *minor* Wahlenb., Fl. Upsal. 33. 1820. [Type: “Hab. in pratis et pascuis fertilibus plerisque frequenter”; *Poa humilis* Ehrh., Beitr. Naturk. 6: 84. 1791, nom. nud.; *P. humilis* Ehrh. ex Hoffm., Deutschl. Fl. 1: 45. 1800, type: “In cultis, ad vias; fl. Apr. Sept.”; *P. subcaerulea* Sm., Engl. Bot. 14, lam. 1004. 1802, basion., type: “Gathered in Anglesea by the Rev. H. Davies, flowering in June”; *P. depressa* J. Presl & C. Presl, Fl. Čech. 20. 1819, type: “Summa Sudetorum cacuminal”; *P. pratensis* var. *latifolia* Weihe ex Mert. & W.D.J. Koch, Deutschl. Fl. 1(2): 612. 1823, type: “Auf dürren sandigen Hügeln, auf mageren Grasplätzen und auf den Triften hoher Gebirge bleibt”; *P. pratensis* var. *subcaerulea* (Sm.) Sm., Engl. Fl. 1: 126 (1824); *P. pratensis* var. *humilis* (Ehrh. ex Hoffm.) Ehrh. ex Spenn., Fl. Friburg. 1: 130. 1825; *P. pratensis* subsp. *latifolia* (Weihe ex Mert. & W.D.J. Koch) Schübl. & G. Martens, Fl. Würtemberg 77. 1834; *P. pratensis* var. *depressa* (J. Presl & C. Presl) Opiz, Seznam Rost. Kvet. Cesk. 76. 1852; *P. pratensis* var. *maritima* Corb., Nouv. Fl. Normandie 655. 1894, type: “Sables maritimes et pelouses du littoral. C.”; *P. irrigata* Lindm., Bot. Not. 1905: 73, 88. 1905, type: “Hab. in uliginosis, pratis et viarum marginibus irrigatis, fossis graminosis, solo abiegnorum muscoso humido, haud raro in pratis litoralibus, hinc inde in pascuis solo duriore turfoso. Vidi specimina typica ex Ölandia (Borgholm),”; *P. irrigata* f. *rigens* Lindm., Bot. Not. 1905: 90. 1905, type: “Hab. In Lapponia”; *P. pratensis* “race” *subcaerulea* (Sm.) Rouy, Fl. France 14: 283. 1913; *P. pratensis* subsp. *irrigata* (Lindm.) H. Lindb., Exsicc. (Pl. Finland.) 2: 20. 1916; *P. pratensis* subsp. *subcaerulea* (Sm.) Hiitonen, Suomen Kasvio 205, fig. 5. 1933]. This variety is known from N and C Europe (introduced in N America) and also appears on N and SW portions of the Iberian Peninsula [**Spa.:** B H Na S SS (Z)], where it is found on grasslands, nitrified dunes and mountainous limestone rocks [0–1380 m a.s.l. May to September].

Two varieties with abundant extravaginal shoots and basal-leaf ligules without hairs or with hairs that are smaller than 0.15 mm, are recognised, although plants having intermediate characteristics are also frequently present. *Poa pratensis* var. *pratensis* [*Poa glabra* Ehrh., Beitr. Naturk. 6: 82. 1791, nom. nud.; *P. pratensis* var. *anceps* Gaudin, Agrost. Helv. 1: 215. 1811, type: “Hab. in paludibus torfaceis. Schleicher.

Perennis”; *P. pratensis* subsp. *anceps* (Gaudin) Lej. & Courtois, Comp. Fl. Belg. 82. 1828; *P. pratensis* α *vulgaris* Gaudin, Fl. Helv. 1: 258. 1828, nom. superfl.; *P. anceps* (Gaudin) Hegetschw., Fl. Schweiz 81. 1838, nom. illeg., non *Poa anceps* G. Forst., Fl. Ins. Austr. 8. 1786; *P. angustifolia* subsp. *anceps* (Gaudin) K. Richt., Pl. Eur. 1: 87. 1890; *P. pratensis* “race” *compressiformis* Rouy, Fl. France 14: 283. 1913, type: “HAB. — Prairies ombragées, tourbières. — Ça et là dans l’aire du type”; *P. pratensis* var. *humilis* sensu Coutinho, Fl. Portugal 1: 104. 1939, non *P. pratensis* var. *humilis* (Ehrh. ex Hoffm.) Spenn., Fl. Friburg. 1: 130. 1825, **syn. nov.**] includes plants in which the blade of most basal leaves is 1.2–3.8(–5.5) mm wide, lanceolate or linear-lanceolate, flat or conduplicate, usually delicate, flexible. This variety is distributed in Eurasia, N Africa and Macaronesia (Azores, Madeira and Canary Islands) and is naturalised in N America and Australia. It is widely dispersed on the Iberian Peninsula [(**And.**) **Spa.**: A Ab Al Av B Bu (C) Cc Co CR Cs (Cu) Ge Gr Gu Hu J L Le Lo Lu M Ma Mu Na O Or P (Po) S Sa Sg So SS (T) Te (To) V Va (Vi) Z Za. **Port.**: AAl (BA) (BB) BL DL E Mi TM], where it appears in mountainous areas [650–2400 m a.s.l. (April) May to July (August)].

Finally, *Poa pratensis* var. *angustifolia* (L.) Sm., Fl. Brit. 1: 105. 1800 [*Poa angustifolia* L., Sp. Pl. 67. 1753, basion., type: “Habitat in Europa ad agrorum versuras” (lectotype designated by Soreng 2000, pg. 254: Herb. Linn. No. 87.12!, excluding second culm from left); *P. brizoides* Vill., Hist. Pl. Dauphiné 2: 126. 1787, nom. illeg., non L. fl., Suppl. Pl. 110. 1782, = *Eragrostis capensis*; *P. villarsii* J.F. Gmel., Syst. Nat., ed. 13, 2: 182. 1791, type: “not expressly indicated”; *P. setacea* Hoffm., Deutschl. Fl., ed. 2, 1: 44. 1800, nom. illeg., non Huds., Fl. Angl. 34. 1762; *P. strigosa* Hoffm., Deutschl. Fl., ed. 2, 1: 44. 1800, type: “In siicioribus elatis; fl. Maj. Iun”; *P. pratensis* subsp. *angustifolia* (L.) Lej., Comp. Fl. Belg. 82. 1828; *P. angustifolia* subsp. *brizoides* K. Richt., Pl. Eur. 1: 88. 1890; *P. pratensis* subsp. *atlantis* Maire, Fl. Afrique N. 3: 101. 1955, type: “Bords des ruisselets, prairies irriguées des collines et montagens siliceuses, rare.- M. Grand Atlas, Mont Gourza vers 2800 m (M.)”], includes plants in which the blade of the basal and shoot leaves is 0.5–1.3 mm wide, linear or setaceous, convolute, usually rigid and brittle. Its distribution area extends across Europe, NW of Africa (Morocco), SW and S Asia and Macaronesia (Canary Islands and Madeira) and it is also introduced in N America. It is scattered over much of the territory covered by *Flora iberica* [**And. Port.**: BL (E) Mi TM. **Spa.**: A Ab Al Av B Bu C Cc Co CR Cu Ge Gr Gu Hu J L Le Lo Lu M Ma Mu Na O Or P PM [Mll] S Sa Sg So (T) Te To V Va Vi Z Za], blooming between April and July, from sea level to 2150 (2400) m.

For a representative list of studied materials, see Suppl. material 1.

2. *Poa legionensis* (Lainz) Fern.-Casas & Lainz in Lainz, Contr. Fl. Asturias 83. 1982

Poa pratensis var. *monticola* Merino, Fl. Galicia 3: 337. 1909, **syn. nov.** [Type: “La var. 2.^a cerca de la cumbre de Peña Rubia (Ancares) á unos 1.700 m. s. m.”]. (Type material probably disappeared).

Poa pratensis subsp. *legionensis* Láinz, Bol. Inst. Estud. Asturianos, Supl. Ci. 15: 43. 1970 [basion.]

Poa alpina subsp. *legionensis* (Láinz) Rivas Mart. & al., Veg. Alta Mont. Cantábrica 279. 1984.

III. Ruiz (1991: 31, lam. III); Fig. 2.

Type. “Ut videtur, diffusa per iuga silicea editissima, occidentalia, montium provinciae legionensis (León) et nonnullarum finitimarum. *Holotypus* in herbario meo hispanico boreo-occidentali: iuxta montem Cornón, pr. Lumajo (Villablino, León), ad 1900 m, 1-VII-1959. Insuper specimina legi simillima, per tractum longum satis: in summo Cellón, pr. Arbas (Rodiezmo, León), ad 2000 m, et paulo inferius, ad 1800 m; item, iuxta lacunam celeberrimi Cueto de Arbas, ad 1750 m, supra Leitariegos (Cangas del Narcea, Asturias); denique, paulo infra summam Peña Trevinca, in ditione quidem zamorensi, ad 2075 m.”. [Holotype JBAG-11515-Láinz!].

Flowering. May to July (September).

Ecology. Wet, somewhat nitrified meadows, stony places, swampy areas and psychro-xerophilic pastures (“cervunales”), on granites or slates; 1560–2400 m a.s.l.

Distribution. Endemic to the CN of the Iberian Peninsula. **Spa.:** Av Cc Le Lu O Or (S) Sa (Za). For a representative list of studied materials, see Suppl. material 1.

3. *Poa cenisia* All., Auct. Fl. Pedem. 40. 1789

Poa distichophylla Gaudin, Alpina 3: 39. 1808. [Type: “Sie kommt ziemlich häufig am Ufer der Alpenbäche im Sand vor, Auf dem Lioson, auf dem Bonhomme in Savoyen, auf den Walliser Alpen u.s. w. Š. Bl. im Jul. und Aug.”].

Poa pallens Haller fil. ex Gaudin, Alpina 3: 41. 1808. [Type: “Dieses zierliche Gras findet man auf den Weiden und an grasigten Felsen auf den höheren Bergen; auf dem großen Bernhard; auf Tzermotanaz au-dessus du val de Bagnes; auf dem Bernardin, unweit der Quelle des Hinter-Rheins, u. s. w. Š. Bl. im Jul. un Aug.”].

Brachypodium cenisium (All) P. Beauv., Ess. Agrostol. 101, 155, 174. 1812.

Poa halleridis Roem. & Schult., Syst. Veg. 2: 539. 1817. [Type: “In alpibus Bernensibus. Inn monte Stockhorn. in Valesia”].

Poa cenisia var. *halleri* Rchb., Icon. Fl. Germ. Helv. 1, ed. 2: 50. 1850. [Type: “In lapidosis rupibusque in Tirol, Bavaria, Helvetia, Pedemontio”].

Poa cenisia subsp. *pallens* (Gaudin) Asch. & Graebn., Syn. Mitteleur. Fl. 2(1): 404. 1900.

Poa fontqueri Braun-Blanq., Bull. Soc. Pharm. Montpellier, Comm. SIGMA 87: 220. 1945. [Type: “Hab. In glareosis schistosis vel graniticis regionis subalpinae-alpinae pyrenaeorum orient. satis frequens. Typus: Pic Fontnègre 2650 m.”].

Poa cenisia subsp. *fontqueri* (Braun-Blanq.) Rivas Mart., Fern. Gonz. & Sánchez Mata, Itinera Geobot. 4: 117. 1990.

Poa cenisia var. *fontqueri* (Braun-Blanq.) Portal, Poa France Belgique Suisse 91. 2005

III. Bolòs and Vigo (2001: 384), Portal (2005: 88, 271, sub var. *cenisia*).




Figure 2. *Poa legionensis* (Lainz) Fern.-Casas & Lainz **A** Habit **B** Detail of the apex of the sheath and ligule, in adaxial view (upper leaf) **C** Spikelet **D** Lower glume, in abaxial view **E** Upper glume, in abaxial view **F** Lemma, in abaxial view **G** Palea, in abaxial view **H** Lodicule **I** Sexual verticils **J** Caryopsis, in adaxial view **K** Caryopsis, in abaxial view. Drawn from MA 314663 and MA 508371.

Type. “Secus torrentes sabulosos exsiccatos in monte Cenisio”. (Type material conserved in TO according to Kerguélen 1975, pg. 238).

Flowering. July to September.

Ecology. Grasslands, gravelly areas, rocky slopes and screes on shale, granite or limestone; edaphically indifferent; 1700–2900 m a.s.l.

Distribution. Mountainous areas of C and S Europe. Mountains of N Spain (Pyrenees, Cantabrian Mountains and N Iberian System). **And. Spa.:** Ge Hu L Le Lo (O) P (S) So Z. For a representative list of studied materials, see Suppl. material 1.

4. *Poa chaixii* Vill., Fl. Delph. 7. 1785 [1786]

Poa sylvatica Chaix in Vill., Hist. Pl. Dauphiné 2: 128. 1787, nom. illeg., non *Poa sylvatica* Pollich, Hist. Pl. Palat. 1: 83. 1776.

Poa sudetica Haenke in Jirasek, Beobacht. Reis. Riesengeb. 120. 1791. [Type: “An dem Aupasturze im Thale”].

Poa rubens Moench, Methodus 187. 1794, nom. illeg., non *Poa rubens* Lam., Tabl. Encycl. 1: 184. 1791; = *Eragrostis unioides*).

Poa latifolia Pohl, Tent. Fl. Bohem. 1: 94. 1810, nom. illeg., non *Poa latifolia* (Osbeck) G. Forst., Fl. Ins. Austr. 8. 1786; = *Centotheca lappaceae*).

Poa sudetica var. *rubens* DC. in Lam. & DC., Fl. Franç. éd. 3, 5: 272. 1815. [Type: “Habitat in Haffia inferior”]. *Poa sulcata* Lag., Gen. Sp. Pl. 3. 1816. [Type: “Legi in sylvis tractus Valgrande dicti, non procul á Pajares oppido”]. Typification supposedly carried out by Fernández Casas & Gamarra (1993: 93), but without establishing an effective lectotypification. Lectotype designated here: “*Poa sulcata* Lam. / ~~*Poa sudetica*~~ / Lagasca iter astur / en Valgrande / Julio.” (label manuscr.): MA 209891].

Cynodon sudeticus (Haenke) Raspail, Ann. Sci. Nat. (Paris) 5: 302. 1825.

Poa chaixii var. *rubens* (DC.) Asch. & Graebn., Syn. Mitteleur. Fl. 2: 423. 1900.

Poa haemi F. Herm., Bull. Soc. Bot. Bulgar. 3: 43. 1929. [Type: “Habitat: Central-Balkan prope Tulovo Bulgariae, ubi legit Iv. Mrkvička Sofia, (Herbarium regale)”].

III. Pignatti (1982: 470), Portal (2005: 93, 273).

Type. “In sylvis & pratis alpestribus circa Chaudun prope Vapincum & ad Taillefer”. (Type material probably conserved in GRM, although doubtful according to Kerguélen 1975, pg. 238).

Flowering. June to August.

Ecology. Mountain meadows and grasslands, in brooms, heaths, beeches, oaks and hollies; edaphically indifferent; 1000–2380 (2780) m a.s.l.

Distribution. C and S Europe; naturalised in Finland. N Iberian Peninsula. **And. Spa.:** (Bu) Ge Hu L Le Lo Lu O (Or) P S (So) (Za). For a representative list of studied materials, see Suppl. material 1.

5. *Poa glauca* Vahl in Oeder, Fl. Dan. 6(17): 3. 1790 subsp. *glauca*

Poa caesia Sm., Fl. Brit. 1: 103. 1800. [Type: “Ang. Sea-green Meadow-grafs. In Scotiâ. D. Fairbairn. Mountains in Bredalbane. Mr. Mackay”].

Poa nemoralis var. *glauca* (Vahl) Gaudin, Agrost. Helv. 1: 182. 1811.

Poa nemoralis subsp. *glauca* (Vahl) Gaudin, Fl. Helv. 1: 240. 1828.

Poa balfourii Parnell, Ann. Mag. Nat. Hist., ser. 1: 10. 1842. [Type: “not expressly indicated”].

Paneion glaucum (Vahl) Lunell, Amer. Midl. Naturalist 4: 222. 1915.

III. Portal (2005: 98, 278, sub var. *glauca*).

Type. “Legi tantummodo in paroecia Wang Walders, ad pedes montium, in Finmarkia minus frequens. Prater aliis notis, praesertim colore glau.....”. (Holotype conserved in C according to PAF 2018: Norway: Oppland, Vang, “legi in alpinis Norvegiae Valders versus Vang”, leg. J. Vahl).

Flowering. (July) August to September.

Ecology. Rocky places and forest and scrub grasslands; (1500) 1900–2770 m a.s.l.

Distribution. Circumboreal: Eurasia (extending S to the Pyrenees, S Alps and N of Greece) and Arctic and alpine regions of N America; also Argentina. NE Spain. (**And.**).

Spa.: Ge Hu. For a representative list of studied materials, see Suppl. material 1.

6. *Poa nemoralis* L., Sp. Pl. 69. 1753 subsp. *nemoralis*

Poa angustifolia var. *nemoralis* (L.) Huds., Fl. Angl., ed. 2: 41. 1778.

Paneion nemorale (L.) Lunell, Amer. Midl. Naturalist 4: 222. 1915.

III. Portal (2005: 282).

Type. “Habitat in Europa ad radices montium umbrosas” (lectotype designated by Soreng 2000, pg. 255: icon in Scheuschzer, Agrostogr. Helv. Prodr. t. 2, 1708; epitype designated by Soreng and Edmondson in Soreng 2000, pg. 255: BM).

Flowering. April to August (November).

Ecology. Grasslands in shady, usually deciduous forests and in pastures, margins of alpine wetlands (“borreguiles”) and fissures of rocks; edaphically indifferent; (135) 550–2980 m a.s.l.

Distribution. Europe, temperate Asia and NW Africa (Morocco); introduced in other parts of the world (e.g. Canada, USA, Patagonia and Guatemala). N half and S third of the Iberian Peninsula. **And. Por.:** BA DL TM. **Spa.:** (A) (Ab) Al Av B Bu Ca Cc Cs Cu Ge Gr Gu Hu J L Le Lo Lu M (Ma) Na Or O P S Sa Sg So SS T Te To V Va Vi Z Za.

Notes. *Poa nemoralis* is a polymorphic species with two recognised patterns of variation and numerous transitional forms in the territory encompassed by *Flora iberica*. The first recognised variety, *Poa nemoralis* var. *nemoralis* [*Poa cinerea* Vill., Hist. Pl.

Dauphiné 2: 126. 1787, type: “Il vient au même endroit que le précédent -*Poa angustifolia*-.”; *P. debilis* Thuill., Fl. Env. Paris ed. 2: 43. 1799, type: “Habitat in pratis”; *P. miliacea* DC. in Lam. & DC., Fl. France ed. 3, 3: 64. 1805, type: “.. par M. Ramond, qui l’a trouvée dans les Pyrénées; ...”; *P. nemoralis* var. *montana* Gaudin, Alpina 3: 27. 1808, type: “In den Wäldern des Jura Š. Bl. im Jun. un Jul.”; *P. nemoralis* subsp. *vulgaris* Gaudin, Agrost. Helv. 1: 179. 1811, nom. superfl.; *P. nemoralis* subsp. *firmula* Gaudin, Agrost. Helv. 1: 181. 1811, type: “In plantici dumetis non rara”; *P. nemoralis* subsp. *coarctata* Gaudin, Agrost. Helv. 1: 185. 1811, type: “In aridis apicisque, ad muros etiam alpinis hawd infrequens”; *P. nemoralis* var. *miliacea* (DC.) Godr. in Gren. & Godr., Fl. France 3: 541. 1856; *P. nemoralis* var. *eunemoralis* Hack. in Briq., Prodr. Fl. Corse 1: 141. 1910, nom. inval.], is widely distributed in the area covered by *Flora iberica* [**And. Por.:** BA DL TM. **Spa.:** Al Av B Bu Ca Cc Cs Cu Ge Gr Gu Hu J L Le Lo Lu M Ma Na Or O P S Sa Sg So SS T Te To V Va Vi Z Za]. This taxon includes plants that are usually green or sometimes glaucous, generally with smooth and flexible stems and with most leaves flat and flexible, linear or linear-lanceolate and erect or erect-patent. The other variety, *Poa nemoralis* var. *rigidula* Mert. & W.D.J. Koch, Deutschl. Fl. 1: 617. 1823 [type: “Auf Wiesen, im Gebüsch der Triften, am Saume der Wälder”], is dispersed throughout the region [**And. Port.:** TM. **Spa.:** (A) (Ab) Al Av Bu Cc Gr Hu L M Na P S Sg Te Z] and includes plants that are usually glaucous, with rigid and often rough stems and usually convolute leaves that are setaceous, linear or linear-lanceolate, ± rigid and erect. For a representative list of studied materials, see Suppl. material 1.

7. *Poa compressa* L., Sp. Pl. 69. 1753

Poa planiculmis Weber, Suppl. Fl. Holsat. 3. 1787. [Type: “Habitat in collibus prope Neumuhlen”].

Poa compressa var. *depauperata* Mutel, Fl. Franç. 4: 81. 1837. [Type: “plante étiolée, forêts des Alpes et du Jura”].

Poa compressa var. *langeana* (Rchb.) W.D.J. Koch, Syn. Fl. Germ. Helv. ed. 2: 932. 1844.

Panicum compressum (L.) Lunell, Amer. Midl. Naturalist 4: 222. 1915.

Poa cenisia subsp. *sardoa* E. Schmid, Vierteljahrsschr. Naturf. Ges. Zürich 78: 239. 1933. [Type: “Perdas Crapias am Gennargentu, 1820 m, Gneissfelsflur (22. VII.1923, b.) ...”].

III. Portal (2005: 95, 276), Soreng and Peterson (2012: 11, fig. I, F-M).

Type. “Habitat in Europae & Americae septentrionalis siccis, muris, tedis” (lectotype designated by Soreng 2000, pg. 255: LINN-87.41!).

Flowering. May to September (October).

Ecology. Meadows and grasslands, forests, gravelly areas, margins of roads and slopes; edaphically indifferent, although preferring basic substrates; (75) 540–1990 (2300) m a.s.l.

Distribution. Circumboreal (most of Europe, to SW Asia); introduced in N, C and S America (Peru, Argentina) and Australia. CN and E Spain, rarer in the south.

And. Port.: TM. **Spa.:** A (Ab) B (Bi) Bu (Cs) Cu Ge Gr Gu H Hu J L (Le) Lo (Lu) M Na (Or) P S Sa Sg So T Te (V) Va Vi (Z) Za. For a representative list of studied materials, see Suppl. material 1.

Notes. Plants of this species usually have a glaucous green colour. The lower (and sometimes higher) leaves of many studied herbarium specimens are missing their blades and are frequently fragmented with the ligule exposed. Inflorescences of *P. compressa* are frequently narrow and interrupted, with almost adpressed branches and spikelets are variable in size and number of flowers. The spikelets may be long, almost always entirely glabrous and glaucous, with 4–9 flowers or short and bear 2–5 flowers, and this variability may be present in the same population or even on the same plant. *Poa compressa* has not been previously listed in the flora of Portugal.

8. *Poa laxa* Haenke in J. Jirasek, *Beobacht. Reis. Riesengeb.* 118. 1791 subsp. *laxa*

Poa laxa f. *pallida* Lange, nom. nud., in sched. (C 10022611; COI-Willk. 36552), **syn. nov.**
III. Bolòs and Vigo (2001: 386), Portal (2005: 108, 281).

Type. “[Haenke 1791: 116, Schneekoppe] der kahle, steinigte Gipfel” [= the summit area of Mt Sněžka, NE Bohemia, at the Polish border]” (lectotype designated by Kirschner et al. 2007, pg. 349: *Poa laxa* a me descripta in Actis Societ. Boh. Anno 1787. *Poa* Halleri historia Nr. 1457. Lecta in Sudetis et in Styriae Alpibus, T. Haenke (PR); epitype designated by Kirschner et al. 2007, pg. 349: Bohemia, the Krkonoše Mts, Mt Sněžka, scree site just below the summit plateau at the beginning of the track called Jubilejní cesta, 50°44'10"N, 15°44'25"E, 3 Jul 2007, J. Zahradníková&L. Harčariková s.n: PRA 349; isoeotypes: PR, PRC).

Flowering. July to September.

Ecology. Rocky places, stony places and high mountain waterfalls, on shale, schist and granite; (1900) 2300–3150 m a.s.l.

Distribution. C and N of Europe, reaching the Carpathians, Balkans, Apennines and Pyrenees, and N America. N and NE Iberian Peninsula. **And. Spa.:** Ge Hu L (P). For a representative list of studied materials, see Suppl. material 1.

9. *Poa minor* Gaudin, *Alpina* 3: 44. 1808

Poa laxa subsp. *minor* (Haenke) Hooker fl., Stud. Fl. Brit. Isl., ed. 1: 444. 1870.

Poa laxa var. *minor* (Haenke) Fiori in Fiori & Paoletti, Fl. Italia 1: 86. 1898.

III. Bolòs and Vigo (2001: 387), Portal (2005: 111, 281).

Type. “Dieses schöne Gras findet man wie das vorige, auf hohen Gebirgen; auf dem Bernhard, auf den Bergen oberhalb Ber und Aigle u. s. w. Bl. im Jul. und Aug.”. (Type material conserved in LAU according to Kerguelen 1975, pg. 241).

Flowering. (July) August to September.

Ecology. Talus slopes, stony places and fissures of rocks, wet and sheltered grasslands, on limestone and schist; 2000–3350 m a.s.l.

Distribution. Mountains of S Europe: Sierra Nevada, Pyrenees, Alps, Balkans and Carpathians. N and SE Spain: Cantabrian Mountains, Pyrenees and Sierra Nevada. (**And.**). **Spa.:** Ge Gr Hu (Na) O S.

Two subspecies of *P. minor* are recognised in the territory encompassed by *Flora iberica*. A key to their identification is given below:

- 1 Blades of most leaves conduplicate, rarely flat, those of basal and shoot leaves 0.4–1.6(–2.1) mm wide; branches of the inflorescence flexuous; spikelets ovate-oblong; lemma hairy at the nerves and the base, hairs of the latter usually longer than the width of the lemma; anthers 0.8–1.1 mm **a. subsp. *minor***
- Blades of most leaves flat, rarely conduplicate, those of basal ones 1.5–2.3 mm wide; branches of the inflorescence straight; spikelets oblong; lemma glabrous or only weakly hairy at the nerves; anthers (1–)1.2–1.7 mm
..... **b. subsp. *nevadensis***

a. subsp. *minor*

Poa supina Panz. in Sturm, Deutschl. Fl. 34: 1. 1812, nom. illeg., non *Poa supina* Schrader, Fl. Germ. 289. 1806.

Poa pyrenaica Lange ex Willk. & Lange, Prodr. Fl. Hispan. 1: 80. 1861, nom. inval., pro syn.

Poa jacetana gr. *laxa* P. Montserrat, nom. nud., in sched. (MA 291693), **syn. nov.**

Flowering. (July) August to September.

Ecology. Talus slopes, stony places and fissures of limestone rocks, wet and sheltered high mountain locations; rarely on schist; 2000–3207 m a.s.l.

Distribution. Mountain systems of S Europe (Cantabrian Mountains, Pyrenees, Alps, Balkans and Carpathians). N Spain: Cantabrian Mountains and Pyrenees. (**And.**). **Spa.:** Ge Hu L Na O S. For a representative list of studied materials, see Suppl. material 1.

b. subsp. *nevadensis* Nannf. in Font Quer, Exsicc. Fl. Iber. Select. Cent. 3: n. 201. 1935.

Poa laxa sensu Boiss., Voy. Bot. Espagne 659. 1842, non *Poa laxa* Haenke in J. Jirasek et al., Beobacht. Reis. Riesengeb. 118. 1791.

Poa minor var. *nevadensis* (Nannf.) Á.M. Hern., Acta Bot. Malac. 2: 35. 1976, comb. inval., **syn. nov.**

III. Fig. 3.




Figure 3. *Poa minor* subsp. *nevadensis* Nannf. **A** Habit **B** Detail of the apex of the sheath and ligule, in adaxial view (upper leaf) **C** Spikelet **D** Lower glume, in abaxial view **E** Upper glume, in abaxial view **F** Lemma, in abaxial view **G** Palea, in abaxial view **H** Lodicule **I** Sexual verticils **J** Immature caryopsis, in adaxial view **K** Caryopsis, in abaxial view. Drawn from GDA 31029 and MA 422687.

Type. “Baetica: in schistosis montium Sierra Nevada, l. Cerro de la Alcazaba dicto, ad 3000 m alt., Leg. Font Quer, 28 aug. 1923” (lectotype designated here: “Institut Botanicum Barcinonense / Flora Iberica Selecta / Cent. III Dec. 1935 / 201. *Poa minor* Gaud. / Fl. Helvet., I, p. 253 (1828). / ssp. *nevadensis* Nannf. nov. ssp. / *Poa laxa* Boiss., Voy. Bot. Esp., II, p. 659, non Haenke. / Baetica: in schistosis montium Sierra Nevada, l. Cerro de la Alca- / zaba dicto, ad 3000 m alt. *Cotypus*. / Leg. Font Quer, 28 aug. 1923. / Obs.: Differt a typo foliis tenuioribus, planis; ligulis brevioribus / (2 mm non excedentibus); spiculis angustioribus et longioribus; / floribus distantioribus; glumis valde inaequalibus; gl. I 2-2,5 mm / longis, gl. II 2,2-3,= mm; antheris paulo longioribus et angustiori- / bus (longit. 1,1-1,3 mm). J. A. Nannfeldt.” (label printed): UPS-V-873177, top specimen on the right; isolectotypes designated here: BC 990150, BC 87706, BC 87707; GDA 31029, 31030, 31031; MA 11385; MAF 28595).

Flowering. July to September.

Ecology. Pastures and wet stony places in high mountain locations, on schist; (2500) 3000–3350 m a.s.l.

Distribution. Endemic to SE Spain: Sierra Nevada. **Spa.:** Gr. For a representative list of studied materials, see Suppl. material 1.

10. *Poa trivialis* L., Sp. Pl. 67. 1753 subsp. *trivialis*

Poa dubia Leers, Fl. Herborn. 28, tab. 6, fig. 5. 1775. [Type: “H. in pratis humidiusculis ubique frequens; ad fossas; an der Dillae & Mühlbach copiose; etiam fisscioribus locis gramineis pervulvaris”].

Poa scabra Ehrh., Beitr. Naturk. 6: 83. 1791, nom. inval.

Poa stolonifera Haller ex Muhl., Descr. Gram. 139. 1817, nom. illeg., non *Poa stolonifera* Bellardi, Mem. Reale Accad. Sci. Torino 5: 215. 1792.

Poa trivialis α. *vulgaris* Rchb., Icon. Fl. Germ. Helv. 1: 37, fig. 1653, 1654. 1834, nom. superfl.

Poa feratiana Boiss. & Reut., Pugill. Pl. Afr. Bor. Hispan. 128. 1852. [Type: “Hab. In sylvâ Irati Pyrenaeorum occidentalium loco Erreca-Idorra (Férat in herb. Fauché!) Fl. Julio”] (lectotype designated by Burdet et al. 1981, pg. 577: specimen indicated with arrows, G00176550!).

Poa sylvicola Guss., Enum. Pl. Inarim.: 371, tab. 18. 1854. [Type: “In sylvaticis apricis ubique vulgatissima; nec non prope Neapolina, et Stabias”].

Poa trivialis var. *umbrosa* Balansa, Bull. Soc. Bot. France 21: 16. 1874. [Type: “Rhizè, dans les forêts, les lieux humides”].

Poa trivialis var. *sylvicola* (Guss.) Hack., Verh. K. K. Zool.-Bot. Ges. Wien 1890: 127. 1890.

Poa attica sensu Pérez Lara, Anales Soc. Esp. Hist. Nat. 15: 406 (1886), non *Poa attica*-Boiss. & Heldr. in Boiss., Diagn. Pl. Orient. ser. 1, 13: 57 (1854).

Poa attica var. *gaditana* Pérez Lara ex Willk., Suppl. Prodr. Fl. Hispan. 22. 1893, **syn. nov.** [Type: « ...provinciae Gadit. Haud frequens (pr. Jerez in loco et Albaladejo et in Dehesa del Torongil; pr. Grazalema ad Huertas de Benamahona, PER. LARA!) »].

Poa trivialis var. *flaccida* Willk. ex J.J. Rodr., Fl. Menorca 152. 1904, **syn. nov.** [Type: “Barranco de Son Blanc en sitios frescos”] (lectotype designated here: first label: “Glumelle glabra. Ax. de l’épilles numi / à la base des glumelles inferieures d’un “ faisceau de poile longe et soyeux. / Gaines sublisses. -illegible- noueux à / leur base, lisses au sommet. / = *P. attica* Boiss. Heldr. var. *flaccida*.” (manuscr.); second label: “J. J. Rodriguez. (printed) Plantas de Menorca. (Balears.) (printed) / *Poa trivialis* L.? (manuscr.) / var. (?) *flaccida* Wk. ined (manuscr.) / Localidad (printed) Barranco de Son Blanc (Algar) (manuscr.) / Estacion (printed) Sitios frescos (manuscr.) / Epoca (printed) 28 mayo 1874 (manuscr.) / Leg. Rodríguez (impreso)”: COI-Willk. 36527; isolectotype designated here: HGM 3121-1).

Poa trivialis subsp. *sylicola* (Guss.) H. Lindb. fil., Oefvers. Förh. Finska Vetensk.-Soc. 38(13): 9. 1906.

Poa trivialis “rasse” *majorica* [*majorcica*] F. Hermann, Verh. Bot. Vereins Prov. Brandenburg 54: 252. 1914, **syn. nov.** [Type: “An Rainen und Wegrändern sammelte ...”].

Poa trivialis f. *majorica* (F. Hermann) Knoche, Fl. Balear. 1: 310. 1921, **syn. nov.**

Poa trivialis f. *flaccida* (Willk. ex J.J. Rodr.) Knoche, Fl. Balear. 1: 311. 1921, **syn. nov.**

Poa trivialis var. *obtusata* Maire, Fl. Afr. N. 3: 99. 1955, nom. inval.

Poa trivialis subsp. *feratiana* (Boiss. & Reut.) Á.M. Hern., Acta Bot. Malac. 2: 33. 1976.

Poa portalii H. Scholz, Willdenowia 42: 293 (2012), **syn. nov.** [Type: Holotype: France: Aquitaine, Pyrenees Atlantiques, Iraty, c. 50 m behind camping place on a wet depression at border of a foot path in woodland, 1150 m, 11.8.2010, Böhling 15255 (B, as “*Poa supina*”)]. Holotype B 100558216 (seen by Soreng in 2015, pers. comm.).

Poa trivialis var. *modesta* Caball., nom. nud., in sched. (MA 11644, MA 11627), **syn. nov.**

Poa trivialis f. *biflora* Bernis, Flora Maragata, nom. nud., in sched. (MA 11615; MA 11617), **syn. nov.**

Poa verticillata auct., non L. (MA 11631).

Poa trivialis var. *contracta* Pérez Lara, nom. nud., in sched. (MAF 28658, MAF 28569, MAF 28655), **syn. nov.**

III. Ruiz (1991: 29, lam. II), Devesa (1987: 263, sub *P. trivialis* subsp. *trivialis* and subsp. *sylicola*).

Type. “Habitat in Europae pasenis” (neotype designated by Soreng 2000, pg. 256: Hudson 16, Herb. Linn. No. 87.9!).

Flowering. (March) April to July.

Ecology. Pastures, hygrophilous grasslands (stream edges, peat bogs, reed patches and meadows) and very wet soils of deciduous forests (e.g. alders, ashes, chestnuts and oaks); edaphically indifferent; 0–2000 (2150) m a.s.l.

Distribution. Europe, N Africa, Asia and Macaronesia (Azores, Madeira and Canary Islands); introduced in other parts of the world. Most of the Iberian Peninsula and Balearic Islands. **And. Port:** AAl Ag BA BB BL (BAI) DL E Mi R TM. **Spa.:** A Ab Al Av B Ba Bu C Ca Cc Co CR Cs Ge Gr Gu H Hu J L Le Lo Lu M Ma (Mu) Na O (Or) P PM[Mll Mn] Po S Sa Se Sg So (SS) (T) (Te) To V Va Vi Za (Z). For a representative list of studied materials, see Suppl. material 1.

Notes. *Poa trivialis* is variable with regard to habit, leaf size and inflorescence morphology. The most distinctive feature of this species is the elongated ligule, which is always longer than the width of the leaf blade, ovate or \pm triangular in the basal leaves and irregularly dentate or bilobed with an acute apex in the uppermost ones. In addition, the spikelets have 2 or 3 flowers and the sharp, arched glumes converge around the lemma. The base of the lemma is very woolly or extremely rarely glabrous and the hairs are clearly longer than its width.

Some plants have somewhat thickened and constricted stolons, with a more or less moniliform appearance. These individuals were described as *Poa sylvicola* Guss. (= *P. attica* var. *gaditana* Pérez Lara ex Willk.; = *P. trivialis* var. *umbrosa* Balansa). According to Soreng (pers. comm.), *P. trivialis* subsp. *sylvicola* (Guss.) H. Lindb. fil. is common in the Mediterranean region, while subsp. *trivialis* is rather infrequent and, conversely, subsp. *sylvicola* is infrequent northwards. Other characteristics of subsp. *sylvicola* are smoother sheaths and the consistent presence of hairs on the lower part of the marginal lemma veins vs. their absence in subsp. *trivialis*. Practically speaking, the marginal vein is hairy in Mediterranean populations but glabrous or nearly so in northern ones. In both types of populations, however, the differing combinations of forms of these characters makes it almost impossible to delimit these two taxa. Consequently, we have opted not to recognise them as separate subspecies.

Poa feratiana Boiss. & Reut. is also included here as a synonym of *P. trivialis*. Plants labelled as *P. feratiana* on herbarium sheets had 2 flowers per spikelet, which is diagnostic for this species, but this characteristic is also very common on most studied sheets of *P. trivialis*. In addition—as indicated by Hernández Cardona (1976) after studying the type material (G-herbarium Boissier)—some of the characteristics attributed to this species in the original description were incorrect. For instance, the number of veins of the lemma is actually 5, not 3, a feature likely overlooked by the original authors because the marginal veins are usually very close to the edge. As another example, the lemma is indeed woolly at the base, as is common in *P. trivialis*.

Plants with some of their spikelets completely sterile and reduced to a set of whitish or hyaline membranes are also known.

11. *Poa flaccidula* Boiss. & Reut., Pugill. Pl. Afr. Bor. Hispan. 128. 1852

Poa balearica Porta, Nuov. Giorn. Bot. Ital. 19: 324. 1887. [Type: “M. Ad pedes rupium praeruptarum m. Coma den Ar-bona” (lectotype designated by Rosselló and Sáez 2000, pg. 141, second on left specimen: G).]

Poa trivialis subsp. *balearica* (Porta) Gand., Nov. Consp. Fl. Eur. 506. 1910, **syn. nov.**

Poa trivialis f. *balearica* (Porta) Knoche, Fl. Balear. 310. 1921.

Poa trivialis var. *balearica* (Porta) O. Bolòs & Molinier, Collect. Bot. (Barcelona) 5: tab. 7. 1958, comb. inval., **syn. nov.**

Poa zapateri Gandoger, nom. in sched. (MA 11500). [Type: “Sierra de Albarracín”]

Poa ventalloi Sennen, nom. nud., in sched. (Sennen BC-966297).

III. Devesa (1987: 264).

Type. “Habitat in umbrosis septentrionalibus jugi Cerro de San Cristoval et Sierra de la Nieve ditionis Serrania de Ronda Junio 1849 (Boiss. et Reuter)” (lectotype designated by Hernández Cardona 1978, pg. 104-105, 337, left specimen on the sheet from Cerro de San Cristóbal, Reuter, 6-1849: G 00176652!).

Flowering. April to July.

Ecology. Stony places, cliffs, scrub clearings and understoreys, on limestone; (200) 700–2030 m a.s.l.

Distribution. W Mediterranean region (S France [Provence], peninsular Spain, Balearic Islands and NW Africa [Morocco and Algeria]). S and E half of peninsular Spain and Mallorca. **And. Spa.:** A (Ab) (Al) B Bu Ca (Co) CR Cs Cu Gr Gu Hu J L Ma (Mu) (Na) PM[Mill] (S) So (T) Te To V Z. For a representative list of studied materials, see Suppl. material 1.

Notes. Spikelets in this species usually have 2–3 flowers, but can have 3–7, a rare phenomenon observed more frequently in populations in NE Spain. The most distinctive characteristic of *Poa flaccidula* is the sericeous or appressed-hairy indumentum of the intervein zone of both the lemma and palea. This species is sometimes confused with *P. annua*, but, along with other differences, the latter is an annual, not a perennial. *Poa flaccidula* can also be confused with *P. trivialis*, which, like *P. flaccidula*, has a ligule that is longer than the width of the leaf blade and possesses hairs at the base of the lemma that are longer than its width; however, both the palea and the internerval surface of the lemma is glabrous in *P. trivialis*.

P. flaccidula subsp. *guadianensis* F.M. Vázquez, Folia Bot. Extremad. 9: 66 (2016) has recently been described from Extremadura (SW Spain), but examination of the type material (HSS 65616; COF 62937 isotypus) reveals that this taxon is in no way attributable to *P. flaccidula*. The most we can say, given the immaturity of the specimens, is that it may be of hybrid origin, with *P. bulbosa* possibly one of the parents.

12. *Poa annua* L., Sp. Pl. 68. 1753 subsp. *annua*

Poa annua var. *viridis* Lej. & Courtois, Comp. Fl. Belg. 1: 80. 1828. [Type: “not expressly indicated”].

Poa ovalis Tineo, Pl. Rar. Sicil., fasc. 2: 21. 1846. [Type: “In pascuis montosis, apricis, palustribus. Cotrano al Gurgo lo Drago”].

Poa annua var. *aquatica* Asch., Fl. Brandenburg 1: 844. 1864. [Type: “Provinz Brandenburg S. Altd. Sumpf hinter dem Pfarrgarten”].

Poa annua var. *typica* Beck, Fl. Nieder-Österreich: 84. 1890, nom. inval.

Poa annua var. *ovalis* (Tineo) Trab. in Batt. & Trab., Fl. Algérie (Monocot.) 2: 206. 1895.

Poa annua f. *plicata prostrata* Sennen, Pl. Espagne n. 605. 1908, nom. nud., in sched. (MA 11165), p.p., **syn. nov.**

Poa annua var. *lanuginosa* Sennen, Diagn. Nouv. sér. 1933: 209, n. 8980. 1936. [Type: “Hab.- Maroc: Melilla à Rostrogordo. Leg. Hno. Mauricio”] (lectotype designated here: “1933.–Plantes d’Espagne. –F. Sennen / N° 8980 / *Poa annua*

L. / var. *lanuginosa* Sennen / Maroc: Melilla, à Rostrogordo / 2–III Leg. Hno. MAURICIO” (label printed): specimen upper on the left, BC 119353; isolecto-type: MA 11155).

Poa annua f. *lanuginosa* Sennen & Mauricio, Cat. Fl. Rif Orient.: 132. 1934, nom. nud., **syn. nov.**

Poa annua var. *pilantha* Ronniger, Verh. Deutsch. Bot. Ges. Wien 88-89: 97. 1941. [Type: “not expressly indicated”, but the material was collected on the island of Zante, Ionian Islands, Greece].

P. annua subsp. *pilantha* (Ronniger) H. Scholz, Ber. Deutsch. Bot. Gesell. 81: 19. 1968.

Ochlopoa annua (L.) H. Scholz, Ber. Inst. Landschafts- Pflanzenökologie Univ. Hohenheim, Beih. 16: 58. 2003.

Ochlopoa annua subsp. *pilantha* (Ronniger) H. Scholz & Valdés, Willdenowia 36: 661. 2006.

III. Ruiz (1991: 27, lam. I), Soreng and Peterson (2012: 14, fig. 2A–E), Devesa (1987: 261).

Type. “Habitat in Europa ad vias” (lectotype designated by Soreng 2000, pg. 254: right-hand plant, Herb. LINN No. 87.17!).

Flowering. All year.

Ecology. Pastures and grasslands along roads, fallow fields, gardens, margins of water-courses and more or less nitrified soils of all types; edaphically indifferent; 0–2100 m a.s.l.

Distribution. Cosmopolitan, although apparently of Mediterranean origin. Throughout the Iberian Peninsula and Balearic Islands. **And. Port:** AAl Ag BA BAl BB BL DL E Mi (R) TM. **Spa.:** A Al Av B Ba Bi Bu C Ca Cc Co CR Cs Cu Ge Gr Gu H Hu J L Le Lo Lu M Ma Mu Na O (Or) P PM [Mll, Mn] Po S Sa Se Sg So SS T Te To V Va Vi Z Za. For a representative list of studied materials, see Suppl. material 1.

Notes. Plants are found in the territory covered by *Flora Iberica* that have hairy lemmas, at least towards the internodal basal zone, with this indumentum being more perceptible in apical flowers of the spikelet. This characteristic is usually accompanied by a very dense silky indumentum in the veins. In other cases, the spikelet has lemmas with a glabrous internodal surface, usually accompanied by a lower density of indumentum in the nerves, with sometimes even the medium ones being glabrous or glabrescent. The first variation corresponds to *Poa annua* var. *lanuginosa* Sennen (Diagn. Nouv. sér. 1933: 209, n. 8980. 1936), a name that prevails over the name *P. annua* var. *pilantha* Ronninger (Verh. Deutsch. Bot. Ges. Wien 88-89: 97. 1941). When Scholz in Ber. Deutsch. Bot. Gesell. 81: 19 (1968) raised Ronninger’s taxon to the subspecies category, he stated that the distribution of this subspecies was Mediterranean (e.g. Greece, Italy, Spain and Morocco) and extra-Mediterranean for the type subspecies. Although plants of Mediterranean environments in the Iberian Peninsula tend to have hairier lemmas, we have also found specimens assignable to var. *annua* and, conversely, we have observed plants with hairy lemmas in typically Eurosiberian areas (e.g. Lugo, Minho, Oviedo and Santander) and even Macaronesia (e.g. Madeira).

In certain populations, some spikelets are completely sterile and reduced to a set of whitish or hyaline membranes. Although infrequent, plants with loosely antrorse-scabrid inflorescence branches have been detected, perhaps as a result of hybridisation with other species (e.g. MA 420475, MA 449625).

DNA sequence data support the hypothesis that *P. annua*, a tetraploid species, has arisen by hybridisation—and subsequent polyploidisation—between two Eurasian diploid species, the annual *P. infirma* Kunth and the rhizomatous perennial *P. supina* Schrad. (Soreng et al. 2010; Mao and Huff 2012), as suggested by Nannfeldt (1937).

13. *Poa infirma* Kunth in Humb., Bonpl. & Kuntz, Nov. Gen. Sp. 1: 158. 1816

Megastachya infirma (Kunth) Roem. & Schult., Syst. Veg. 2: 585. 1817.

Eragrostis infirma (Kunth) Steud., Nomencl. Bot. ed. 2, 1: 563. 1840.

Poa annua var. *exilis* Tomm. ex Freyn, Verh. K. K. Zool.-Bot. Ges. Wien 27: 469. 1878. [Type: “So auf sonnigen, trockenen Grasplätzen der Macchien, meist in Gesellschaft von Asterolinon, Euphorbia peploides, E. exigua und anderen Zwergpflanzen stellenweise häufig, bisher aber nur läng der Küste von Fasana bis Medolino; auch auf S. Marina (Tommasini 1872)”].

Poa annua var. *remotiflora* Hack. ex Batt. & Trab., Fl. Algérie Monocot. 206. 1895. [Type: “Lieux humides et région montagneuse, Rouïba, Teniet-el-Haàd”].

Poa annua subsp. *exilis* (Tomm. ex Freyn) Murb. in Asch. & Graebn., Syn. Mitteleur. Fl. 2: 389. 1900.

Poa remotiflora (Hack. ex Batt. & Trab.) Murb., Acta Univ. Lund. 36 Afd. 2. n. 1: 22. 1900, nom. illeg., non *Poa remotiflora* Rupr., Fl. Samoied. Cisural. 63. 1845.

Poa exilis (Tomm. ex Freyn) Murb. ex Nannf., Acta Univ. Lund., ser. 2, 1(4): 73. 1906.

Poa annua f. *plicata prostrata* Sennen, Pl. Espagne n. 605. 1908, nom. nud., in sched. (MA 11165), p.p., **syn. nov.**

Poa annua var. *plicata* Sennen, Pl. d’Espagne w.n. 1908, nom. nud., in sched. (MA 11142).

Poa annua var. *laxiflora* Sennen, Pl. d’Espagne n. 606. 1908, nom. nud., in sched. (MA 11143).

Poa annua subsp. *remotiflora* (Hack. ex Batt. & Trab.) Jansen & Wachter, Fl. Nederl. 1(2): 78. 1951, nom. illeg.

Poa annua L. var. *spiciformis* Palau Ferrer, Pl. Baleares n. 793. 1955, nom. in sched. (GDA 30993; MA 168790), **syn. nov.**

Ochlopoa infirma (Kunth) H. Scholz, Ber. Inst. Landschafts-Pflanzenökologie Univ. Hohenheim Beih. 16: 59. 2003.

III. Ruiz (1991: 27, lam. I), Soreng & Peterson (2012: 14, fig. 2F–H), Devesa (1987: 262).

Type. “Crescit in frigidis regni Novogranatensis, inter Fontibon, Suba et Santa Fe de Bogota, alt. 1360 hexap. Floret Augusto”. [Holotypus P-HUMB; isotypus: B-WILLD-1974 pl. 223, LE-TRIN-2638.01 fragm. & illustr., US-1851276 fragm. ex P, US-2851277 fragm. ex P-HUMB) (designated by Soreng and Peterson 2012, pg. 43)].

Flowering. October to May (July).

Ecology. Therophytic pastures and ruderal places, preferably in sandy soils; edaphically indifferent; 2–1000 m a.s.l.

Distribution. W Europe, Mediterranean, Macaronesian and Irano-Turanian regions extending to India; introduced in Australia and the Americas. Scattered across the Iberian Peninsula and Balearic Islands. **Port:** AAl Ag BA BAl E Mi (R) TM. **Spa.:** A (Ab) Al Av B Ba (C) Ca Cc Co CR Ge Gr H Hu J (L) Lo M Ma (Mu) O (Or) (Po) PM [Mll Mn] S Sa Se Sg Te To V Va (Z) Za. For a representative list of studied materials, see Suppl. material 1.

Notes. This species is clearly differentiated from the previous one, not only by the small size of its anthers, but also by the smaller size of the leaves, which, in most cases, are not more than 2 mm wide and have margins that are barely scabrid or even smooth. Most populations comprise plants of small size (up to 22 cm).

Plants with some spikelets completely sterile and rudimentary, whitish or hyaline have been detected in some populations (e.g. province of Granada, GDA 15557).

14. *Poa supina* Schrad., *Fl. Germ.* 1: 289. 1806

Poa annua var. *varia* Gaudin, *Alpina* 3: 29. 1808. [Type: “An den Bächen, auf den höheren Apen; auf der Scheideck. Seringe. Auf dem Gotthard häufig. Ø. Bl. im Sommer”].

Poa annua subsp. *varia* (Gaudin) Gaudin, *Agrost. Helv.* 1: 189. 1811.

Poa annua var. *supina* (Schrad.) Spenn., *Fl. Friburg.* 1: 127. 1825.

Poa annua subsp. *supina* (Schrad.) Husn., *Graminées* 51. 1898.

Poa exigua Foucaud & Mandon ex Husn., *Gram. Fr. Belg.* 88. 1899, nom. illeg., non

Poa exigua Dumort., *Gramin. Belg.* 113. 1824.

Poa annua f. *macranthera* Lit. & Maire in Jahand. & Maire, *Cat. Pl. Maroc* 1: 66. 1931. [Type: “Sierra Nevada. Pyrénées. Corse. Alpes.”].

Poa supina f. *exigua* Gamisans, *Candollea* 29: 48. 1974. [Type: “Massif du Cinto, Capo al Berdato, versans SSW, pozzine de pente, 2320 m, 1.8.1969, Gamisans 2912 (fl.); cirque de Trimbolacciu, couloir de Pampanosa, pelouse, 1880 m, 6.8.1970, Gamisans 2913 (fl.)”].

Ochlopoa supina (Schrad.) H. Scholz & Valdés, *Willdenowia* 36: 662. 2006.

III. Pignatti (1982: 469).

Type. “In summis alpiis Salisburgensibus (Mielichhoffer)”. (Possible isotype conserved in LE according to PAF 2018).

Flowering. June to August (September).

Ecology. Perennial grasslands in wet places; edaphically indifferent; 1200–3481 m a.s.l.

Distribution. C and SW Europe, Apennines, Fennoscandia and NE Russia, extending to the Rif (Morocco). Pyrenees, Cantabrian Mountains, Central System and Sierra Nevada. **And. Port.:** (BA)? (Mi)? (TM)? **Spa.:** Av Ge Gr Hu L Le M Na P S. For a representative list of studied materials, see Suppl. material 1.

Notes. Plants from the Sierra Nevada tend to have glabrescent floral parts (lemmas and paleas) or an indumentum that is restricted to the basal zone of the central nerve; this contrasts with plants from some peripheral populations of this mountain massif (e.g. Lugros, Dehesa del Camarate, 2200 m, GDAC 41005) and other Iberian populations, none of which are usually glabrous. Although clearly corresponding to *P. supina*, the Sierra Nevada plants also resemble those of *P. rivulorum* Maire & Trab., Bull. Soc. Hist. Nat. Afrique N. 15: 395. 1924; *P. annua* var. *rivulorum* (Maire & Trab.) Lit. & Maire in Jahand. & Maire, Cat. Pl. Maroc 1: 66. 1931; *P. alpina* subsp. *atlantica* (Trab.) Romo, Treb. Inst. Bot. Barcelona 11: 40. 1987, **syn. nov.**, non *P. alpina* var. *atlantica* Trabut in Maire, Mém. Soc. Sci. Nat. Maroc 7: 147. 1924; *Ochlopoa rivulorum* (Maire & Trab.) H. Scholz & Valdés, Willdenowia 36: 662. 2006], a tetraploid species ($n = 14$) endemic to Alto, Medio and Anti Atlas (Morocco), in that the flowers, as indicated by Maire (1955), are usually glabrous and rarely hairy at the base of the medial and marginal veins. The observed pattern in Sierra Nevada is perhaps simply infrapopulational variation.

15. *Poa bulbosa* L., Sp. Pl. 70. 1753 subsp. *bulbosa*

Poa bulbosa subsp. *eu-bulbosa* Hayek, Prodr. Fl. Penins. Balcan. 3: 259. 1932, nom. inval.

III. Ruiz (1991: 31, lam. III), Devesa (1987: 265).

Type. “Habitat in Gallia” (lectotype designated by Meikle 1985, pg. 1742; restricted by Soreng 2000, pg. 255: Hasselquist, Herb. Linn. No. 87.57!).

Flowering. January to July (December).

Ecology. Pastures and grasslands in wet and nitrified soils, less frequently in ephemeral pastures and dry places; edaphically indifferent; 0–3100 m a.s.l.

Distribution. Europe, SW, C and N Asia until W China, Africa and Macaronesia (Madeira and Canary Islands); introduced in the Americas, Australia and Pacific Islands. Entire Iberian Peninsula and Balearic Islands. **And. Port.:** Ag AAl BAl BA BB BL DL E Mi R TM. **Spa.:** A Al Ab Av B Ba Bu C Ca Cc (Cs) CR Co Cu Ge Gr Gu H Hu J L Le Lo Lu M Ma Mu Na Or O P PM[Mll (Me)] Po S Sa Sg Se So SS T Te To V Va Vi Z Za. For a representative list of studied materials, see Suppl. material 1.

Notes. According to Bolòs and Vigo (2001: 380), reports of *Poa bulbosa* subsp. *concinna* (Gaudin) Hayek, Repert. Spec. Nov. Regni Veg. Beih. 30(3): 260. 1932 [*P. concinna* Gaudin, Agrost. Helv. 1: 196. 1811, basion., type: “Hab. in arenosis Valesiae inferioris, praecipue Seduni”; *P. perconcinna* J.R. Edmonson, Bot. J. Linn. Soc. 76: 330. 1978, type: based on *P. concinna* Gaudin] in E Spain are mistaken, as this taxon is actually only distributed from SE France to the Balkan Peninsula. That taxon differs from *P. bulbosa* by their smaller sizes and narrower leaves, 0.8–2.2 mm ligules and never-proliferating spikelets with 6–10 flowers. In a few peninsular populations, plants with spikelets bearing 9–10 flowers have been detected, but they coincide with *P. bulbosa* in all other characters.

Caryopses are formed in this species, but sexual reproduction is infrequent; more common propagation routes include the formation of bulbs at the base of the plant or bulbils at the inflorescence level. This latter phenomenon, pseudovivipary, is extraordinarily frequent in *P. bulbosa* and involves the formation of bulbils for vegetative multiplication in the place of normal flowers (“proliferated spikelets”; Ofir and Kigel 2014). These bulbils may or may not coexist with normal flowers on the same spikelet or plant or in the same population. The balance between clonal and sexual reproduction is controlled mainly by day length and temperature; short days and low temperatures usually promote proliferating inflorescences, whereas long days and high temperatures induce normal and seminiferous ones. The proliferating spikelets usually carry (1–) 2–3 bulbils and have deformed floral parts: the glumes are usually narrower, the lemma is typically long (up to 20 mm), thin and either glabrous or only hairy on the central and marginal veins and the palea is missing or fully integrated into the bulbil, similar to the lodicules. Stamens are also missing or very reduced in size. The bulbils tolerate desiccation, are dormant during the summer and are dispersed by wind and ants. Both the basal bulbs, which are also dormant during the summer and the inflorescence bulbils sprout at the peak of the winter rainy season (cf. Ofir and Kigel 2014).

Two varieties are distinguished in the flora. Plants with bulbils in the inflorescence are recognised as *Poa bulbosa* var. *vivipara* Koeler, Descr. Gram. 189. 1802 [Type: “prope Moguntiam in arenosis”; *Poa bulbosa* subsp. *vivipara* (Koeler) Arcang., Comp. Fl. Ital. 785. 1882; *Paneion bulbosum* var. *viviparum* (Koeler) Lunell, Amer. Midl. Naturalist 4: 222. 1915; *Poa bulbosa* f. *vivipara* (Koeler) Maire, Fl. Afrique N. 3: 86. 1955], a variety distributed in Europe, Africa and SW, C and N Asia to W China and introduced in the Americas, Australia and the Pacific Islands; it appears in practically all of the provinces of *Flora iberica* [**And. Port.:** AAl BAl BB BL E TM. **Spa.:** A Al Ab Av B Ba Bu Ca Cc CR Co Cu Ge Gr Gu H Hu J L Le Lo Lu M Ma Mu Na Or P PM[Mll] S Sa Sg Se So SS Te To V Va Vi Z Za], from sea level to 3100 m, with a preference for shady places. The other variety, *Poa bulbosa* var. *bulbosa* [*Poa crispa* Thuill., Fl. Env. Paris, ed. 2: 45. 1799, type: “Habitat in locis arenosis. Floret Maio”; *P. pasqualii* Heldr. ex Parl., Fl. Ital. 1: 343. 1850, nom. inval., pro syn.; *P. bulbosa* subsp. *perligulata* H. Scholz, Bot. Chron. 3: 17. 1983, Typus: “Italia: Insula Elba, marina di Campo, 12.4.1980, W. Lang w.n. (B)”]; *P. perligularis* H. Scholz, Willdenowia 16: 404. 1987, non *Poa perligulata* Pilger, Notizbl. Bot. Gart. Berlin-Dahlem 11: 779. 1933; *P. bulbosa* f. *variegata* Font Quer, nom. nud., in sched. (MA 11272), **syn. nov.**; *P. bulbosa* f. *minor* H. Villar, nom. nud., in sched. (MA 156764, MA 156765), **syn. nov.**], is distributed throughout the range of the species and comprises plants in which the spikelets produce normal flowers; this variety is found mostly in provinces covered by *Flora iberica* [**Port.:** Ag AAl BAl BA BB BL DL E Mi R TM. **Spa.:** A Al Ab Av (B) Ba (Bi) Bu C Ca Cc (Cs) CR Co Cu Ge Gr Gu H Hu J L Le Lo Lu M Ma (Na) Or O P PM[(Ib) Mll (Me)] Po S Sa Sg Se So (SS) T Te To V Va Vi Z Za], where it ranges from sea level to 2895 m and generally thrives in exposed places.

Bulbs appear to have arisen in *Poa* at least twice and possibly as many as four times (Cabi et al. 2016). Taxa having bulbs are distributed in four clades: (1) *P. supersect. Poa*

(*P. densa* Troitsky and *P. diversifolia* (Boiss. & Balansa) Hack. ex Boiss; (2) in or near *P. supersect.* *Homalopoa* (*P. pseudobulbosa* Bor); (3) N-clade (*P. pelasgis* H. Scholz); and (4) *P.* subgen. *Ochlopoa* sect. *Arenariae* (type: *P. bulbosa*) mixed with species of *P.* sect. *Alpinae* (type: *P. alpina*).

In Sierra de Mariola (Alicante), a population has been detected (MA 752991) in which the ligule of the upper leaves in some individuals is very short (1–1.5 mm), subtruncate and irregularly dentate or not.

16. *Poa ligulata* Boiss., Voy. Bot. Espagne 2: 659. 1842

Poa concinna var. *membranacea* Boiss., Elench. Pl. Nov. 89. 1838, nom. subst. (Burdet et al. 1981 stated that they had not seen material corresponding to this taxon in herbarium G; however, they forget that this name has been replaced by *P. ligulata* and, therefore, the type material is the same).

Poa djurdjurae Trab. in Batt. & Trab., Fl. Alger 207. 1884. [Type: “Col de Tirourda (juin 1883)”].

Poa paui Font Quer, Iter Marocc. n. 34. 1928, nom. in sched. [Type: “Hab. in glareosis calc. montis Tisuka (Gomara), 2100 m. / alt.; 13 junii.” (lectotype designated here: “Dr. Font Quer. – Iter Maroccanum, 1928 / 34. *Poa paui* F. Q., sp. nov. / Hab. in glareosis calc. montis Tisuka (Gomara), 2100 m. / alt.; 13 junii. / Descr.: *Poa ligulata* affinis, sed folia subquadruplo lon-/ giora, ligulis subduplo elongatis et angustioribus; panicula laxa, / glumis virescentibus, spiculis majoribus.” (label printed): BC 70616, lower central specimen; isoelectotypes designated here: BC 70616a, b, and c; MA 11378; GDA 31039)].

Poa ligulata var. *paui* (Font Quer) Maire, Bull. Soc. Sci. Nat. Maroc 11: 113. 1931.

Poa ligulata var. *djurdjurae* (Trab.) Maire, Bull. Soc. Hist. Nat. Afrique N. 22: 323. 1931.

Poa membranacea (Boiss.) C. Vicioso, Anales Jard. Bot. Madrid 2: 192. 1942, nom. illeg.; Anales Jard. Bot. Madrid, 6(2): 13. 1946.

Poa ligulata var. *eu-ligulata* Maire & Weiller in Maire, Fl. Afrique N. 3: 87. 1955, nom. inval.

Poa ligulata var. *mauretanica* Maire, Fl. Afrique N. 3: 88. 1955. [Type: “M. Beni Snassen au Ras Foughal! (E.); Moyen Atlas ! (M., J., E.) ; Rif (F.-Q., M.)”].

III. Bolòs and Vigo (2001: 378), Devesa (1987: 265), Fig. 4.

Type. [Loc. ind. “Hab. in glareosis frigidis in sumâ Sierra Tejeda et in Sierra Nevada loco Corral dicto”] (lectotype designated by Hernández Cardona 1978, pg. 267, 358: top specimen on the right, G 00418689!, Herb.-Boiss.; isoelectotypes: G 00418689a, b, cl, Herb.-Boiss.; ex herbier Reuter-Barbey, G 00418691!, 00418691a!; ex herbier De Candolle, G00418690!; vide Burdet et al. 1981, pg. 578).

Flowering. (April) May to July.

Ecology. Rocky places, pastures in protosols and stony places, on limestone and dolomite, less frequently on schist or gypsum; (700) 740–3200 m a.s.l.




Figure 4. *Poa ligulata* Boiss. **A** Habit **B** Detail of the apex of the sheath and ligule, in adaxial view (upper leaf) **C** Spikelet **D** Lower glume, in abaxial view **E** Upper glume, in abaxial view **F** Lemma, in abaxial view **G** Palea, in abaxial view **H** Lodicule **I** Sexual verticils **J** Caryopsis, in adaxial view **K** Caryopsis, in abaxial view. Drawn from MA 414266 and MA 423181.

Distribution. Iberian Peninsula and NW Africa. E half of Spain, rarer towards the W. **Spa.:** (A) (Ab) Al Bu Ca Cc Cs CR Cu Gr Gu J Le Lo M Ma Mu Na (Or) P S Sg Se So Te To V Z. For a representative list of studied materials, see Suppl. material 1.

Notes. This species is unmistakable because of its large ligules, which are very exerted, decurrent on the leaf sheath—especially those of the base and shoots—and pearly white, which makes its white-green tufts very striking. Its presence is indicated for the first time in the region of Extremadura (Spain).

17. *Poa alpina* L., Sp. Pl. 67. 1753 subsp. *alpina*

III. Soreng and Peterson (2012: 11, fig. I, A–E), Pignatti (1981: 473).

Type. “Habitat in alpinis Lapponicis, Helveticis” (lectotype designated by Soreng 2000, pg. 254: Herb. Linn. No. 87.2!).

Flowering. June to August (September).

Ecology. Pastures in pine forests, fir woods, beeches and bushes, ruderalised hills and rocky plains, on limestone, schist, granite and slate; 1200–3150 m a.s.l.

Distribution. Circumboreal: Europe, Asia, N America, locally at low elevations in NW Africa (Morocco). N Spain, Cantabrian Mountains, Pyrenees and the Iberian System. **And. Esp.:** (Av) B Bi Ge Hu L Le Lo Na O P S (Sa) So Te Vi. For a representative list of studied materials, see Suppl. material 1.

Notes. A characteristic of this species is the size and shape difference of the ligule on the basal leaves and shoots—especially the oldest ones—compared with that of the upper leaves. In the first case, the ligule is tiny, with a more or less complete margin; in the second case, the ligule is noticeably larger, often with an irregular margin and sometimes even split into two or more parts.

This is a very polymorphic taxon, with four recognised, sometimes intergrading varieties in the territory encompassed by our revision. Plant size, leaf stiffness and panicle contraction can vary extensively depending on altitude, exposure and soil type. The first of these varieties is *Poa alpina* var. *alpina* [*Poa frigida* Gaudin, *Alpina* 3: 33. 1808, type: “au-dessous du glacier de Plan-nové; dans la vallée de Bagnes etc.”; *P. alpina* var. *frigida* (Gaudin) Salisb., *Flora* 16(2): 473. 1833; *P. alpina* var. *genuina* Godr. in Gren. & Godr., *Fl. France* 3: 543. 1855, nom. inval.; *P. alpina* var. *involucrata* Lange, *Pugill. Pl. Hispan.* 47. 1860, **syn. nov.**, type: “In regione alpina Pyren. Hisp. ad Port de Benasque (9 Aug. c. fl.)”, lectotype designated here: “Herb. Joh. Lange (printed) / *Poa alpina* var. *involucrata* nob. / ligulio foliorum omnium truncatis, ver. / -illegible- infer. folio -illegible- tetra / Port de Benasque reg. alp. 9 Aug. 1851. (manuscr.)”: C 10022547, specimen on the left), Fig. 5; *P. nuriensis alpina* Sennen, *Bull. Soc. Bot. France* 73: 677. 1926, nom. nud., **syn. nov.**; *P. alpina* subsp. *digitata* Beauverd, *Bull. Soc. Bot. Genève, sér. 2*, 26: 122, fig. 2a–g. 1936, type: “Hab. in locis apricis calidisque valliculae dictae “du Grand Tabuc” ad locum dic-



Figure 5. Lectotype of *Poa alpina* var. *involucrata* Lange (C 10022547). Reproduced with permission of the Natural History Museum of Denmark.

tum “les Grangettes”, ca. 1800 m alt., supra thermis “Le Monétier de Briançon”, 15 Julii 1933, leg. J. Vergnet et G. Beauverd”; *P. badensis* subsp. *multiflora* sensu Rivas. Mart., Itinera Geobot. 15: 705. 2002, non *P. alpina* var. *multiflora* Gaudin, Fl. Helv. 1: 245. 1828; *P. nuriensis* Sennen, Pl. Espagne n. 4063. 1916, nom. nud., in sched. (BC 70515; MA 11298), **syn. nov.**]. This circumboreal taxon includes large plants having large, generally delicate, non-rigid leaves with non-thickened margins—or thickened less than 0.05 mm—and developed inflorescences that are only slightly condensed. This variety is typically found in sheltered, less-exposed locations and is widespread throughout the range of the species [**And. Spa.:** (Av) B Bi Ge Hu L Le Lo Na O P S (Sa) So Te Vi].

Under more adverse conditions, the plants are usually small and possess short, stiff leaves with non-thickened, or up to 0.05 mm thick, margins and very contracted panicles. These plants have been designated as *Poa alpina* var. *brevifolia* (Gaudin) Godr. in Gren. & Godr., Fl. France 3: 543. 1855 [*Poa alpina* subsp. *brevifolia* Gaudin, Fl. Helv. 1: 245. 1828, basion., type: “in M. Sempronio ad pylas vallis Ganter”], a variety distributed in mountains of C Europe and extended through the central and eastern Pyrenees [**And. Spa.:** B Hu L]. This pattern of variation is probably clinal and needs to be checked experimentally. The third recognised variety, *Poa alpina* var. *molinerii* (Balb.) Endl., Cat. Horti Vindob. 46. 1842 [*Poa molinerii* Balb., Elenco 85. 1801, basion., type: “Locis saxosis, et siccis prope Tenda reperta est, ac in hortum Taurinensem adlata ab eximio Ignatio MOLINERI, cujus triviale nomen imposui, utpote ejus stirpis inventore”], comprises plants having leaves with whitish, cartilaginous, thickened (0.1–0.15 mm) margins that, together with the middle underside vein, form a clear, visible contrast to the green leaf blade, the latter mostly flat or conduplicate and rigid. This taxon, is distributed in mountains of S and C Europe and, to date, only two populations of *P. alpina* var. *molinerii* have been detected for *Flora iberica*, one in **Andorra** (Coll de Ordino, on the way to Casamanya, 2100 m; MA 514862) and the other in **Spain** (Lérida, Clot del Muniyidor, 2215 m; BC 877255).

Finally, *Poa alpina* var. *vivipara* L., Sp. Pl. 67. 1753. [Type: “An haec α . β . sequentis tantum varietas; Lectotype designated by Soreng 2000, pg. 254: three left-hand culms, LINN-87.4!; *Poa vivipara* (L.) Willd., Enum. Pl. 103. 1809; *P. alpina* subsp. *vivipara* (L.) Arcang., Comp. Fl. Ital. 785. 1882; *P. alpina* f. *vivipara* (L.) B. Boivin, Naturaliste Canad. 94: 628. 1967], includes plants with pseudoviviparous spikelets. This variety, distributed mainly in N and C Europe, Greenland, Iceland and N America, has only been detected in a population in the territory covered by *Flora iberica* (**Spain:** Na), namely, it was collected in the valley of Roncal at 1600 m a.s.l (SEV 97164). The inflorescences of these plants conserved on this sheet show a great contrast of colours: glumes and lemmas of light green and straight or curved, dark green proliferations.

Another taxon described for the flora of Morocco is *Poa alpina* subsp. *stenobotrya* Maire (Bull. Soc. Hist. Nat. Afrique N. 33: 95. 1942), which is distinguished by its linear-lanceolate panicle and the presence of hairs at the base of the lemmas.

Conclusions

In the territory covered by *Flora iberica*, the genus *Poa* is represented by 24 taxa (17 species, 1 subspecies and 8 varieties), mostly perennial. The majority of these taxa have broad global and/or European distributions, whereas two (*P. legionensis* and *P. minor* subsp. *nevadensis*) are Spanish endemics and two have restricted distributions (*P. ligulata*, Iberian–North African; *P. flaccidula*, Iberian–North African and the Balearic Islands, extending to Provence, France). The most widely distributed species are *P. bulbosa* and *P. annua*, reflecting their worldwide range. The provinces with the greatest representation of *Poa* are Huesca, Santander, Lérida and Andorra, all located in the N Iberian Peninsula, which are traversed by mountain systems and subjected to a temperate climate.

Acknowledgements

We are grateful to the curators of BC, C, COI, GDA-GDAC, HSS, JACA, MA, MAF, MGC, SALA-SALAF, SEV and UNEX for arranging loans, A. Herrero (MA), N. Ibáñez and J. Pallàs (BC), O. Ryding (C-Lange), L. Loze (G-Boiss.), O. Garcia (HGM), J.I. Alonso (JBAG), J. Pizarro (MAF) and M. Hjertson (UPS) for sending photos, R. Morales (Royal Botanical Garden of Madrid) for translating the type protologues published in German Gothic script by some authors and R. Pino (LOU) for his help in trying to locate Merino type material. The study was supported by the Spanish Ministry of Science and Innovation through Projects CGL2008-0298-Co3-3, CGL2011-28613-C03-02 and CGL2014-52787-C8-3-P, all co-financed by the European Regional Development Fund. The English text was edited and improved by Dr. B. Goodson (Edanz Group, www.edanzediting.com/ac). Finally, we thank M. Olonova (Tomsk State Univ.) for her suggestions and express special gratitude to R. Soreng (Smithsonian Inst.) for his excellent comments that improved our paper.

References

- Bolòs O, Vigo J (2001) *Flora dels Països Catalans*, vol. 4. Editorial Barcino, Barcelona, 1–750.
- Burdet HM, Charpin A, Jacquemoud F (1981) Types nomenclatureaux des taxa ibériques décrits par Boissier ou Reuter. I. Gymnospermes à Graminées. *Candollea* 36: 543–584. http://bibdigital.rjb.csic.es/HTML/Burdet_Types_I_Gymnospermae-Gramineae.pdf
- Cabi E, Soreng RJ, Gillespie L, Amiri N (2016) *Poa densa* (Poaceae), an overlooked Turkish steppe grass, and the evolution of bulbs in *Poa*. *Willdenowia* 46(2): 201–211. <https://doi.org/10.3372/wi.46.46201>
- Devesa JA (1987) *Poa* L. In: Valdés B, Talavera S, Galiano EF (Eds) *Flora de Andalucía Occidental*, vol. 3. Editorial Ketres, Barcelona, 260–265.
- Edmonson JR (1980) *Poa* L. In: Tutin TG, Heywood VH, Burges NA, Moore NA, Valentine DH, Walters SM (Eds) *Flora Europaea*, vol. 5. Cambridge University Press, Cambridge, 159–167.

- Fernández Casas J, Gamarra R (1993) Herbarium Lagascanum. Fontqueria 36: 67–108. <http://bibdigital.rjb.csic.es/spa/Volumenes.php?Libro=2699>
- Flora iberica (2018) <http://www.floraiberica.es/> [accessed 22.03.2018]
- Franco JA, Rocha ML (1998) Nova Flora de Portugal (continente e Açores), vol. 3. Escolar editor, Lisboa, 1–283.
- Gillespie LJ, Soreng RJ, Bull RD, Jacobs SWL, Refulio-Rodriguez NF (2008) Phylogenetics relationships in subtribe Poinae (Poaceae, Poeae) based on nuclear ITS and plastid *trnT-trnL-trnF* sequences. Botany 86: 938–967. <https://doi.org/10.1139/B08-076>
- Giussani LM, Gillespie LJ, Scatagliini MA, Negritto MA, Anton AM, Soreng RJ (2016) Breeding system diversification and evolution in American *Poa* supersect. *Homalopoa* (Poaceae: Poeae: Poinae). Annals of Botany 118(2): 281–303. <https://doi.org/10.1093/aob/mcw108>
- Hernández Cardona AM (1976) Notas sobre el género *Poa* en la Península Ibérica. Acta Botanica Malacitana 2: 31–38. <http://www.biolveg.uma.es/abm/abm.html>
- Hernández Cardona AM (1978) Estudio monográfico de los géneros *Poa* y *Bellardiochloa* en la Península Ibérica e islas Baleares. Dissertationes Botanicae 46: 1–365.
- Kerguelen M (1975) Les gramineae (Poaceae) de la Flore Française. Essai de mise au point taxonomique et nomenclaturale. Lejeunia 75: 1–343.
- Kirschner J, Kirschnerová L, Štěpánek J (2007) Generally accepted plant names based on material from the Czech Republic and published in 1753–1820. Preslia 79: 323–365.
- Maire R (1955) *Poa* L. Flore de l'Afrique du Nord, vol. 3. P. Lechevalier, Paris, 76–101.
- Mao Q, Huff R (2012) The evolutionary origin of *Poa annua*. Crop Science 52(4): 1910–1922. <https://doi.org/10.2135/cropsci2012.01.0016>
- Meikle RD (1985) Flora of Cyprus, vol. 2. The Bentham-Moxon Trust, Royal Botanic Gardens, Kew, 1741–1746.
- Nannfeldt JA (1937) The chromosome numbers of *Poa* sect. *Ochlopoa* A. & Gr. and their taxonomical significance. Botaniska Notiser 1937: 238–254.
- Ofir M, Kigel J (2014) Temporal and intraclonal variation of flowering and pseudovivipary in *Poa bulbosa*. Annals of Botany 113(7): 1249–1256. <https://doi.org/10.1093/aob/mcu037>
- PAF (2018) Panarctic Flora. Annotated checklist of the Panarctic Flora. Vascular plants. <https://nhm2.uio.no/paf/>
- Pignatti S (1982) Flora d'Italia, vol. 3. Edagricoltore, Bologna, 1–780.
- Portal R (2005) *Poa* de France, Belgique et Suisse. Ed. Vals près, R. Portal, Baltimore, Md., 1–303.
- Rosselló JA, Sáez LI (2000) Index Balearicum: an annotated check-list of the vascular plants described from the Balearic Islands. Collectanea Botanica (Barcelona) 25(1): 3–203. <http://collectaneabotanica.revistas.csic.es/index.php/collectaneabotanica/article/viewArticle/42>
- Ruiz T (1991) *Poa* L. In: Devesa JA (Ed) Las graminéas de Extremadura. Servicio de Publicaciones de la Universidad de Extremadura, Badajoz, 23–33. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=5767>
- Soreng RJ (2000) *Poa*. In Cafferty S, Jarvis CE, Turland NJ (Eds) Typification of Linnaean plant names in the Poaceae (Gramineae). Taxon 49(2): 239–260.
- Soreng RJ, Barrie FR (1999) Proposal to conserve the name *Poa pratensis* (Gramineae) with a conserved type. Taxon 48(1): 157–159. <https://doi.org/10.2307/1224640>

- Soreng RJ, Bull RD, Gillespie LJ (2010) Phylogeny and reticulation in *Poa* based on plastid *trnTLF* and nrITS sequences with attention to diploids. In: Seberg O, Petersen G, Barford AS, Davis JI (Eds) Diversity, phylogeny, and evolution in the Monocotyledons. Aarhus University Press, Aarhus, 619–644.
- Soreng RJ, Gillespie LJ, Consaul LL (2017) Taxonomy of the *Poa laxa* group, including two new taxa from Arctic Canada and Greenland, and Oregon, and a re-examination of *P. sect. Oreinos* (Poaceae). *Nordic Journal of Botany* 35(5): 513–538. <https://doi.org/10.1111/njb.01507>
- Soreng RJ, Peterson PM (2012) Revision of *Poa* L. (Poaceae, Pooideae, Poeae, Poinae) in Mexico: New records, re-evaluation of *P. ruprechtii*, and two new species, *P. palmeri* and *P. wendtii*. *PhytoKeys* 15(0): 1–104. <https://doi.org/10.3897/phytokeys.15.3084>
- Soreng RJ, Peterson PM, Romaschenko K, Davidse G, Zuloaga FO, Judziewicz EJ, Filgueiras TS, Davis JI, Morrone O (2015) A worldwide phylogenetic classification of the Poaceae (Gramineae). *Journal of Systematics and Evolution* 53(2): 117–137. <https://doi.org/10.1111/jse.12150>
- Soreng RJ, Peterson PM, Romaschenko K, Davidse G, Teisher JK, Clark LG, Barberá P, Gillespie LJ, Zuloaga FO (2017) A worldwide phylogenetic classification of the Poaceae (Gramineae) II: An update and comparison of two 2015 classifications. *Journal of Systematics and Evolution* 55(4): 259–290. <https://doi.org/10.1111/jse.12262>
- Stoneberg Holt SD, Horová L, Bureš P (2004) Indel patterns of the plastid DNA *trnL-trnF* region within the genus *Poa* (Poaceae). *Journal of Plant Research* 117(5): 393–407. <https://doi.org/10.1007/s10265-004-0172-4>
- Thiers B [continuously updated] Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/> [accessed 20.03.2018]
- Watson L, Dallwitz MJ (1992) The Grass genera of the World. Cab International, Wallingford, 718–719.
- Willkomm M (1870) *Poa* L. In: Willkomm M, Lange J (Eds) *Prodromus Florae Hispanicae*, vol. 1. Stuttgartiae, 80–83.
- Willkomm M (1893) *Supplementum Prodromi Florae Hispanicae*. Stuttgartiae, 1–370.

Supplementary material I

List of representative sheets for each province and taxon, excluding type material

Authors: Ana Ortega-Olivencia, Juan A. Devesa

Data type: species data

Explanation note: The total number of studied sheets is also indicated for each taxon.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.103.26029.suppl1>